



Peter D. McCann, MD  
Director, Orthopaedic Surgery and  
Orthopaedic Strategic Initiatives,  
Lenox Health Greenwich Village  
Pmccann1@northwell.edu  
www.shouldersurgerynyc.com



Northwell Health Physician Partners  
Orthopaedic Institute at LHGV  
7 7<sup>th</sup> Avenue, 2nd Floor  
New York, NY 10011  
(646) 665-6784

## Elbow Exercises

### Elbow Stretch


A. With the opposite hand, hold the fingers of the affected arm up and stretch for 5 seconds as illustrated. Rest for several seconds, then repeat the stretch for 10 repetitions.

B. Perform a similar stretch, but this time then the wrist of the affected arm down and hold and stretch for 5 seconds. Rest for several seconds and repeat for 10 repetitions.



### Twist

Hold a towel with two hands and twist the towel to the maximum position with the hand in the opposite direction. Hold the twisted position for 5 seconds, then relaxed. Repeat for 10 repetitions.



### Elbow Flexion

Hold a 5 pound weight in the hand of the affected arm and fully bend the elbow. Performed three sets of 10 repetitions and then increase weight in the hand as comfort permits.


Peter D. McCann, MD  
Director, Orthopaedic Surgery and  
Orthopaedic Strategic Initiatives,  
Lenox Health Greenwich Village  
Pmccann1@northwell.edu  
www.shouldersurgerynyc.com

Northwell Health Physician Partners  
Orthopaedic Institute at LHGV  
7 7<sup>th</sup> Avenue, 2nd Floor  
New York, NY 10011  
(646) 665-6784


## Elbow Exercises



**Forearm Rotation**  
Hold a 2 pound weight in the hand of the affected arm and slowly rotate the forearm palm up, then palm down. Performed three sets of 10 repetitions and then increase weight as comfort permits.



**Wrist Curl**  
In the hand of the affected arm and with the forearm facing you, hold a 2 pound weight and bend the wrist slowly up. Hold for 5 seconds, then rest. Repeat for three sets of 10 repetitions. Increase weight as comfort permits.



**Wrist Extension**  
In the hand of the affected arm, hold a 2 pound weight with the forearm and hand facing downwards. Slowly bend the wrist up and hold for 5 seconds then rest. Repeat for three sets of 10 repetitions. Increase weight as comfort permits.